On-line Data Supplement

The Clinical and Pathologic Features of Familial Interstitial Pneumonia (FIP)

Mark P. Steele, Marcy C. Speer, James E. Loyd, Kevin K. Brown, Aretha Herron, Susan H. Slifer, Lauranell H. Burch, Momen M. Wahidi, John A. Phillips III, Thomas A. Sporn, H. Page McAdams, Marvin I. Schwarz, and David A. Schwartz

Methods-Supplement


ATS Dyspnea Scale: In this scale, dyspnea is rated as 0-absent; 1-shortness of breath when hurrying on level ground or up a slight hill; 2-walking slower than people your own age on the level because of breathlessness; 3-having to stop for breath when walking at your own pace on the level; 4-having to stop for breath after walking 100 yards (or after a few minutes) on the level; or 5- too breathless to leave the house or breathless on dressing or undressing.


Supplemental Figure Legend:


Supplemental Figures E1-E7. Pedigree figures depict relationships and diagnostic status for 111 FIP pedigrees. Not all individuals with possible FIP, unaffected, or unknown individuals are shown on pedigrees as a result of pedigree trimming for presentation purposes.

PF Definite NSIP Definite NSIP Probable PF Probable Other ILD Definite PF Possible Other ILD Probable Indeterminate Unaffected **COP Definite** ? RBILD Probable NA RBILD Possible

Supplemental Figure E1







Supplemental Figure E2

